



# AMERICAN FRONTLINE NURSES

## IVERMECTIN WEIGHT CONVERSIONS FOR THE I-CARE EARLY INTERVENTION PROTOCOL

HOW OFTEN TO TAKE IVERMECTIN		TWICE A WEEK SPACED OUT	TAKE DOSE DAILY FOR AT LEAST 5 DAYS OR UNTIL SYMPTOMS RESOLVE	TAKE DOSE DAILY FOR AT LEAST 5 DAYS OR UNTIL SYMPTOMS RESOLVE
BODY WEIGHT POUNDS TO KILOGRAMS		PREVENTION DOSING 0.2MG/KG	EARLY INTERVENTION LOW DOSING 0.4MG/KG	EARLY INTERVENTION HIGH DOSING 0.6MG/KG
<b>70-90 LB</b>	32-40 KG	8mg	16mg	24mg
<b>91-110 LB</b>	41-50 KG	10mg	20mg	30mg
<b>111-130 LB</b>	51-59 KG	12mg	24mg	36mg
<b>131-150 LB</b>	60-68 KG	13.5mg	27mg	40.5mg
<b>151-170 LB</b>	69-77 KG	15mg	30mg	45mg
<b>171-190 LB</b>	78-86 KG	16mg	32mg	48mg
<b>191-210 LB</b>	87-95 KG	18mg	36mg	54mg
<b>211-230 LB</b>	96-104 KG	20mg	40mg	60mg
<b>231-250 LB</b>	105-113 KG	22mg	44mg	66mg
<b>251-270 LB</b>	114-122 KG	24mg	48mg	72mg
<b>271-290 LB</b>	123-131 KG	26mg	52mg	78mg
<b>291-310 LB</b>	132-140 KG	28mg	56mg	84mg

**NOTES:**